



UNIVERSITY OF
SURREY



Royal Surrey
NHS Foundation Trust

Pancreatic exocrine insufficiency and pancreatic enzyme replacement therapy

Mary Phillips BSc (Hons) RD DipADP

Advanced Specialist Dietitian (Hepato-pancreatico-biliary surgery)

Royal Surrey County Hospital, UK

Post Graduate Researcher, University of Surrey, UK



WORLD CLASS CARE
FOR OUR COMMUNITY

Declaration of interests: Honoria received for speaking from Mylan, Sanofi, Vitaflo, Nutricia Clinical Care, Abbott Nutrition and Merck.

Introduction: setting the scene



WORLD CLASS CARE
FOR OUR COMMUNITY



BUM PANCREAS

Introduction

Anatomy and
function of the
pancreas

Causes and
incidence of
pancreatic exocrine
insufficiency

Impact of
pancreatic exocrine
insufficiency

Managing
pancreatic enzyme
replacement
therapy

Recommendations
for practice



Poll

What is PEI?

- 1) When the pancreas stops digesting fat
- 2) When the pancreas stops making insulin
- 3) When the pancreas is withered / atrophic
- 4) When the pancreas stops making enough enzymes



WORLD CLASS CARE
FOR OUR COMMUNITY

Definition

“...deficiency or absence of digestive enzymes leading to maldigestion of food and consequently malabsorption of nutrients” (Whitcomb et al, 2010)


“Exocrine pancreatic insufficiency results from a progressive loss of acinar pancreatic cells which leads to the secretion of an insufficient amount of digestive enzyme into the duodenum.” (Pezzilli et al, 2013)

“...pancreas is unable to deliver sufficient amounts of digestive enzymes to the small intestine, leading to maldigestion” (Sikkens et al, 2012)

Failure of the pancreas to secrete sufficient enzymes to achieve normal digestion



Anatomy and function



- Oblong gland 12.5 x 2.5cm
- Consists of endocrine cells: islets of langerhans) which produce glucagon, insulin etc (1% of all cells)
- 99% cells – exocrine function – producing pancreatic enzymes and fluid. (1200-1500mls/day)



Digestive enzymes

Site	Carbohydrate	Fat	Protein
Saliva	Amylase	Salivary lipase	
Gastric Secretion	Gastric Amylase	Gastric Lipase	Pepsin; Rennin; Gelatinase;
Pancreatic Secretion	Amylase	Lipase; Steapsin	Trypsin; Elastase; Chymotrypsin; Carboxypeptidase;
Jejunal / Ileal Secretion	Sucrase; Maltase; Lactase Isomaltase;	Intestinal Lipase	Brush Border Peptidases



Anatomy and
function of the
pancreas

Causes and
incidence of
pancreatic exocrine
insufficiency

Impact of
pancreatic exocrine
insufficiency

Managing
pancreatic enzyme
replacement
therapy

Recommendations
for practice



Causes of PEI

- Lack of Healthy Pancreatic Tissue (Primary Insufficiency)
 - Pancreatic Resection - Chronic Pancreatitis
 - Recovering Acute Pancreatitis - Pancreatic Cancer
 - Pancreatic Trauma - Cystic Fibrosis
- Lack of Pancreatic Stimulation (Secondary Insufficiency)
 - Gastric Resection
 - Duodenal Resection
- Some work suggests insufficiency in Coeliac Disease, Diabetes, Irritable bowel syndrome, intensive care, inflammatory bowel disease and the elderly



Poll

Which of these is NOT correct?

1. Up to 98% of patients will have PEI after a Whipples
2. More than 50% of people with pancreatic cancer will have PEI on diagnosis
3. People with PEI will always have diarrhoea
4. PEI usually gets worse with time



How common is PEI in pancreatic cancer?

- Present in the vast majority of people with pancreatic cancer
- Progressive
- 66-94% of patients have PEI at first presentation (all comers)
- Function deteriorates at approximately 10% per month
- Function tests can take 2-6 weeks to give results
- Incidence after surgery depends on the type of operation
 - 20-80% tail of pancreas (distal pancreatectomy)
 - Up to 98% head of pancreas (pancreatico-duodenectomy / Whipple)

Sikkens et al, 2014, Tseng et al, 2016, Phillips et al, 2021, Phillips M, 2015, Okano, 2016



Pancreatic Function Tests


Faecal Elastase

- <200ug/g moderate PEI
- <100ug/g severe PEI
- 200 – 500ug/g (low sensitivity/specificity)
- >500ug/g: Consider age; Dilutional samples (watery / large volume stool); Sample collection technique

Breath tests

Calibre of pancreatic tissue on imaging: Pancreatic ductal dilatation





WORLD CLASS CARE
FOR OUR COMMUNITY

Poll

Which of these is NOT a symptom of PEI?

- 1) Bloating
- 2) Blood sugars that are lower than usual
- 3) Vitamin and mineral deficiencies
- 4) Failure to gain weight despite eating really well
- 5) New diagnosis of diabetes



Clinical symptoms (1)

Steatorrhoea

- Loose watery yellow/orange stool
- Floats / difficult to flush away
- Oily / visible food particles

LIMITATIONS

- NOT PRESENT in low fat diet
- MASKED by constipating drugs
- VERY LATE symptom



Clinical symptoms (2)

- Large volume stool
 - Undigested food in the stool
 - Post-prandial abdominal pain
 - Nausea / colicky abdominal pain
 - Gastro-oesophageal reflux symptoms
 - Bloating / flatulence
 - *Weight loss despite good oral intake*
 - Vitamin deficiencies (especially A,D,E,K,)
 - *Hypoglycaemia in patients with diabetes*
- (O'Keefe et al, 2001, Genova Diagnostics, 2008, Friess & Michalski, 2009)




Diagnosis

Pancreatic pathology



Clinical symptoms




Likely PEI

Pancreatic pathology



Diagnostic test



Likely PEI

Clinical symptoms



Diagnostic test



Consider PEI and investigate for pancreatic pathology



Anatomy and
function of the
pancreas

Causes and
incidence of
pancreatic exocrine
insufficiency

Impact of
pancreatic exocrine
insufficiency

Managing
pancreatic enzyme
replacement
therapy

Recommendations
for practice



Malignant disease

Survival

- RCT (unresectable ca pancreas)– no benefit; but predominantly tail of pancreas disease (Woo et al, 2016)
- Independently associated with survival in unresectable disease (189 vs. 95 days, retrospective) (Dominguez-Munoz et al, 2018)
- ESPAC studies show the benefit of completing the full chemotherapy regime – performance status....






ELSEVIER

Contents lists available at ScienceDirect

Pancreatology

journal homepage: www.elsevier.com/locate/pan



Enzyme replacement improves survival among patients with pancreatic cancer: Results of a population based study

K.J. Roberts^{a, *}, C.A. Bannister^b, H. Schrem^c

^a Honorary Reader and Consultant Surgeon, Institute of Immunology and Immunotherapy, University of Birmingham, UK

^b Digital Health Laboratories, UK

^c Consultant Surgeon, Dept Visceral, General and Transplant Surgery, Hannover Medical School, Germany



WORLD CLASS CARE
FOR OUR COMMUNITY

Trial design

Retrospective observational study

PDAC 1998 – 2015

Excluded those with concurrent pancreatitis, PERT prior to diagnosis




WORLD CLASS CARE
FOR OUR COMMUNITY

All patients

PERT

Non PERT




WORLD CLASS CARE
FOR OUR COMMUNITY

No surgery, no chemo


PERT

Non PERT



WORLD CLASS CARE
FOR OUR COMMUNITY

Factors influencing survival





Study Overview



RICOCHET 1: A National Prospective Audit of the Diagnostic Pathway for Suspected Pancreatic Cancer



<50% patients on PERT

Ricochet study (2021)

- 45% of unresectable patients prescribed PERT
- 74.4% potentially resectable patients prescribed PERT
- 96.9% of pancreatic head resection patients prescribed PERT
- PERT prescription was more likely if:
 - Seen by a dietitian ($p = 0.001$)
 - Seen in a specialist centre ($p = 0.049$ – HPB; $p = 0.009$ – pancreas)
 - Seen by a clinical nurse specialist ($p = 0.028$)



Implications

- 37.5 % of readmissions after pancreatic surgery caused by malnutrition and dehydration (Grewal et al, 2011)
- Sarcopenia independently associated with PEI (Shintakuya et al, 2017)
- “difficulty with digestion” is most common symptom in long term (Cloyd et al, 2017)
- PEI guidance primary unmet need in pancreatic cancer (Gooden & White, 2013)



Sarcopenia and outcome.....



Applied nutritional investigation

A high visceral adipose tissue-to-skeletal muscle ratio as a determinant of major complications after pancreatoduodenectomy for cancer

Marta Sandini M.D.^a, Davide P. Bernasconi Ph.D.^b, Davide Fior M.D.^c, Matilde Molinelli M.D.^a, Davide Ippolito M.D.^c, Luca Nespoli M.D.^a, Riccardo Caccialanza M.D.^d, Luca Gianotti M.D., Sc.D.^{a,*}




Influence of cachexia and sarcopenia on survival in pancreatic ductal adenocarcinoma: A systematic review

I. Ozola Zalite^a, R. Zyklus^b, M. Francisco Gonzalez^c, F. Saygili^d, A. Pukitis^{a,e}, S. Gaujoux^{f,g}, R.M. Charnley^h, V. Lyadov^{i,*}



Impact of Sarcopenic Obesity on Failure to Rescue from Major Complications Following Pancreaticoduodenectomy for Cancer: Results from a Multicenter Study

Nicolò Pecorelli, MD¹, Giovanni Capretti, MD², Marta Sandini, MD³, Anna Damascelli, MD⁴, Giulia Cristel, MD⁴, Francesco De Cobelli, MD⁴, Luca Gianotti, MD, ScD³, Alessandro Zerbi, MD², and Marco Braga, MD¹



1605-012-1923-5.



Impact of Sarcopenia on Outcomes Following Resection of Pancreatic Adenocarcinoma

Muscle Index and Pancreatic Fistula

Development After Pancreaticoduodenectomy

HIROAKI YAMANE¹, TOMOYUKI ABE¹, HIRONOBU AMANO^{1,2}, KEIJI HANADA³, TOMOYUKI MINAMI³, TSUYOSHI KOBAYASHI², TOSHIKATSU FUKUDA⁴, SHUJI YONEHARA⁵, MASAHIRO NAKAHARA¹, HIDEKI OH DAN² and TOSHIO NORIYUKI^{1,2}



WORLD CLASS CARE
FOR OUR COMMUNITY

Anatomy and
function of the
pancreas

Causes and
incidence of
pancreatic exocrine
insufficiency

Impact of
pancreatic exocrine
insufficiency

Managing
pancreatic enzyme
replacement
therapy

Recommendations
for practice



UK management

Pancreatic enzyme replacement therapy

- Multiple disease aetiology
- Co-morbidities
- Altered dietary intakes
- Altered meal patterns
- Healthy eating vs. nutritional support



Poll

What is a suitable starting dose for PERT?

- 1) 5,000 units with meals
- 2) 10,000 units with meals
- 3) 22-25,000 units with meals
- 4) 44-50,000 units with meals




How do the products compare?

	Amylase	Protease	Lipase
Creon micro (100mg)	3600	200	5000
Pancrex V capsule	9000	460	8000
Creon 10,000	8000	600	10000
Nutrizym 22	19800	1100	22000
Creon 25,000	18000	1000	25000
Pancrease HL	22500	1250	25000
Pancrex V powder (1g)	30000	1400	25000
Creon 40,000	DISCONTINUED		



Dosing



Comparison of weight-based doses of enteric-coated microtablet enzyme preparations in patient with cystic fibrosis

N = 21

Population: Cystic Fibrosis

Open label crossover: 500u/kg with meals and 250U/kg with snacks compared to 1500u/kg with meals and 750u/kg with snacks.

Diet : 100g fat / day

CFA: increased from 86% to 91% (P<0.05)

(Beker et al, J.Paed Gastrol Nutr. 1994 Aug;19(2):191-7).



WORLD CLASS CARE
FOR OUR COMMUNITY

RESEARCH ARTICLE

Clinical validation of an evidence-based method to adjust Pancreatic Enzyme Replacement Therapy through a prospective interventional study in paediatric patients with Cystic Fibrosis

Joaquim Calvo-Lerma^{1,2*}, Jessie Hulst³, Mieke Boon⁴, Carla Colombo⁵, Etna Masip¹, Mar Ruperto⁶, Victoria Fornés-Ferrer¹, Els van der Wiel³, Ine Claes⁴, Maria Garriga⁶, Maria Roca¹, Paula Crespo-Escobar¹, Anna Bulfamante⁵, Sandra Woodcock³, Sandra Martínez-Barona¹, Ana Andrés², Kris de Boeck⁴, Carmen Ribes-Koninckx¹, on behalf of MyCyFAPP project[†]

PLOS ONE | <https://doi.org/10.1371/journal.pone.0213216> March 12, 2019


NHS

Royal Surrey
NHS Foundation Trust

- Multicentre trial
- Cystic fibrosis cohort
- Diet – 40% Lipid; 40% CHO and 20% Protein
- 1622 – 2573kcal/day



WORLD CLASS CARE
FOR OUR COMMUNITY



$$IOD = \frac{g(90\%_{\text{clinical target CFA)}) - \beta_0 - (\beta_1 \cdot \text{transit time}) - (\beta_3 \cdot \text{age}) - (\beta_4 \cdot \text{PPI intake})}{\beta_2}$$

Table 3. Beta regression model to assess the influence of the study variables on CFA, including the dose of enzymes (TOD) and the individual factors intake of proton pump inhibitors (PPI), age and transit time.

	(exp)Estimate	Confidence Interval CI 95%	p-value
(Intercept) (β_0)	2.839	[0.223, 36.147]	0.42
TOD (β_2)	0.999	[0.998, 1.000]	0.13
PPI intake (β_4)	1.367	[0.885, 2.115]	0.09
Age (β_3)	1.013	[0.961, 1.069]	0.62
Transit time (β_1)	1.815	[1.177, 2.797]	0.007

Transit time played a significant role in results:
longer the transit time the greater the CFA



And it is not just fat....

Medium-Chain Triglyceride Absorption in Patients with Pancreatic Insufficiency

S. CALIARI, L. BENINI, C. SEMBENINI, B. GREGORI, V. CARNIELLI & I. VANTINI
Division of Gastroenterologic Rehabilitation, University of Verona, Verona, and Dept. of Pediatrics,
University of Padua, Padua, Italy

4 way Crossover trial: LCT vs. MCT +/- 50,000 units lipase
N= 6, all male Chronic Pancreatitis patients
All had severe exocrine insufficiency (CFA < 80%)
(1 x distal panc, 1 x total gastrectomy, 1 x distal gastrectomy,
1x whipple, 2 on insulin.)






Fig. 2. Mean fecal fat losses (see Fig. 1).




Fig. 4. Mean fecal nitrogen losses (see Fig. 1).



Dosing used in clinical trials

Study	Cohort	Dose	Benefit
Kim et al, Clin Gastroenterol Hepatol. 2019	RCT n=304 Pancreatico-duodenectomy	40,000 units lipase with meals	Increase body weight; increased pre-albumin
Sato et al, Pancreas . 2018 Aug;47(7):800-806	N=88 PDAC chemotherapy	48,000 units lipase with meals	No difference in nutritional markers in 8/52 trial Survival 19/12 vs. 12/12 (p=0.07)
Woo et al, Pancreatology . 2016 Nov - Dec;16(6):1099-1105	N= 67 Unresectable PDAC	25,000 capsules x 6-9 per day	NO difference in nutritional markers or QOL in 8/52 trial
Bruno et al, Gut . 1998 Jan;42(1):92-6	N = 21 Unresectable PDAC	50,000 units lipase with meals; 25,000 units with snacks	12% improvement in CFA; weight gain in intervention; weight loss in placebo



Recommended dose

STARTING DOSE....

44 - 50,000 units with meals

22 - 25,000 units with snacks

25 - 50,000 units with supplements

Will need higher dose with larger meals

Increase until symptom control



WORLD CLASS CARE
FOR OUR COMMUNITY

Poll

Which of these do you NOT need to take PERT with?

- 1) Cup of tea and two bourbon biscuits
- 2) Mug of peppermint tea
- 3) Milkshake
- 4) Cup of milky coffee



Anatomy and
function of the
pancreas

Causes and
incidence of
pancreatic exocrine
insufficiency

Impact of
pancreatic exocrine
insufficiency

Managing
pancreatic enzyme
replacement
therapy

Recommendations
for practice



WORLD CLASS CARE
FOR OUR COMMUNITY

Pancreatic cancer in adults: diagnosis and management

NICE guideline

Published: 7 February 2018

[nice.org.uk/guidance/ng85](https://www.nice.org.uk/guidance/ng85)

1.6 Nutritional management

- 1.6.1 Offer enteric-coated pancreatin for people with unresectable pancreatic cancer.
- 1.6.2 Consider enteric-coated pancreatin before and after pancreatic cancer resection.



Recommendations for clinical practice

CONSENT



Timing

- Mix with food
- Allow for slow meals / multiple courses / gastric emptying

Dose

- Minimum starting dose 50,000u with meals; 25,000u with snacks
- Increase until symptom control
- Snacks vs. meals
- Nutritional supplements

Prevent denaturation




- <25°C
- ?Proton pump inhibitor?
- Avoid swallowing with hot food/fluids



WORLD CLASS CARE
FOR OUR COMMUNITY

BMJ
Open
Gastroenterology

Consensus for the management of pancreatic exocrine insufficiency: UK practical guidelines

Mary E Phillips ¹ Andrew D Hopper,² John S Leeds ³ Keith J Roberts ⁴
Laura McGeeney,⁵ Sinead N Duggan,⁶ Rajesh Kumar⁷

Contra-indications

For Pancrease HL[®]

Should not be used in children aged 15 years or less with cystic fibrosis

For Nutrizym 22[®] gastro-resistant capsules

Should not be used in children aged 15 years or less with cystic fibrosis

Cautions

Can irritate the perioral skin and buccal mucosa if retained in the mouth; excessive doses can cause perianal irritation

Side-effects

Common or very common

Abdominal distension; constipation; nausea; vomiting

Uncommon

Skin reactions

Frequency not known

Fibrosing colonopathy



Contraindications and side effects

- **CONTRAINDICATIONS**

- CONSENT: Porcine content
- Pork allergy / previous intolerance

- **SIDE EFFECTS**

- Nausea
- Gout (uric acid)
- Fibrosing colonopathy

- **PREGNANCY & BREASTFEEDING:**

- Essential fatty acids are needed for brain and retinol development in the first 8 weeks of pregnancy – DO NOT STOP PERT



What do you **NEED** to know :PEI

- Exocrine insufficiency is progressive and doses escalate with time
- Some patients need really high doses (>150,000 units with a meal = >25 capsules / day = 9-10 x 100 cap tubs per month)
- Significant pill burden
- Micronutrient deficiency common
- Enzymes denatured by excess temperature and acid
- Treat like insulin – different doses for different patients for different meals



Conclusion

- PEI is under recognised
- Many patients are on sub-optimal doses
- Appropriate therapy improves outcome
- Multiple factors play a role in dose adjustment – individual management
- Permission to dose escalate
- Other conditions can mimic PEI
- More data needed to explore relationship with survival in pancreatic cancer.



Case 1



WORLD CLASS CARE
FOR OUR COMMUNITY

- 78 year old female
- Metastatic pancreatic cancer – liver mets
- Declined palliative chemotherapy
- Bowels open once a day: type 1 (pellet type stool)
- Large volume hard stool
- Crampy abdominal pain
- 5% weight loss each month despite eating well
- Functionally well – caring for Grandchildren



Poll

Is this patient at risk / potentially suffering from malabsorption?

1. Yes – treat
2. Yes – test and then consider treatment
3. No
4. Not sure



- Commence laxatives
- Once bowel symptoms improved – then start PERT
- Biggest symptom was constipation



Case 2

- 69 year old female
- Localised pancreatic ductal adenocarcinoma
- Distal pancreatectomy and splenectomy – 1cm lesion removed from tail of pancreas
- Bowels open 4 x a day loose and watery
- Gaining weight, 1-2kg per month
- Eating well



Poll

Is this patient at risk / potentially suffering from malabsorption?

1. Yes – treat
2. Yes – test and then consider treatment
3. No
4. Not sure



- Investigate:
 - Total specimen 2cm tail of pancreas
 - Head and body of pancreas enhances normally, no duct dilatation
 - Faecal elastase >500ug/g
 - Vitamin and mineral levels normal



Poll

Is this patient at risk / potentially suffering from malabsorption?

1. Yes – treat
2. Yes – test and then consider treatment
3. No
4. Not sure



Outcome

- No nutritional compromise
- **PEI = Failure of the pancreas to secrete enough enzymes to achieve normal digestion**

PDAC is a Red Herring

- Further investigations – diagnosed small intestinal bacterial overgrowth
- Symptoms resolved with antibiotics.

